



ARL-TR-7706 • JULY 2016



Synthesis of 1,3-Dinitrohexahydropyrimidine via Ring Contraction of Ether-Linked Nitramines

**by Joseph E Banning, William M Sherrill, Eric C Johnson, and
Jesse J Sabatini**

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.



Synthesis of 1,3-Dinitrohexahydropyrimidine via Ring Contraction of Ether-Linked Nitramines

by Joseph E Banning, William M Sherrill, and Jesse J Sabatini
Weapons and Materials Research Directorate, ARL

Eric C Johnson
Bowhead Science and Technology, Belcamp, Maryland

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) July 2016		2. REPORT TYPE Final		3. DATES COVERED (From - To) Fiscal Year 2013	
4. TITLE AND SUBTITLE Synthesis of 1,3-Dinitrohexahydropyrimidine via Ring Contraction of Ether-Linked Nitramines				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Joseph E Banning, William M Sherrill, Eric C Johnson, and Jesse J Sabatini				5d. PROJECT NUMBER AH43	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Research Laboratory ATTN: RDRL-WML-C Aberdeen Proving Ground, MD 21005-5069				8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-7706	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Two ether-linked nitramine explosives have been synthesized. The reactivity of ether-linked nitramines has been probed and it was found that with heating, it is possible to perform a ring contraction to a methylene-linked nitramine that may be suitable for melt-cast applications.					
15. SUBJECT TERMS TNT, melt cast, ring contraction, DHP, ether-linked nitramines					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 20	19a. NAME OF RESPONSIBLE PERSON Joseph E Banning
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 410-278-9656

Contents

List of Figures	iv
List of Tables	iv
Acknowledgments	v
1. Introduction	1
2. Computational Analysis and Sensitivity	2
3. Synthesis Discussion	3
4. Results and Experimental Section	3
4.1 Synthetic Procedures	4
5. Discussion of DHP and Formaldehyde Extrusion from Ether-Linked Nitramines	6
6. Conclusions	7
7. References	9
List of Symbols, Abbreviations, and Acronyms	10
Distribution List	12

List of Figures

Fig. 1	Structures of Compounds 1–3.....	1
Fig. 2	Ring contraction.....	2
Fig. 3	Synthesis of Compound 1	3
Fig. 4	Synthesis of Compound 2	3
Fig. 5	1,3-dinitrotetrahydropyrimidin-2(1H)-one	4
Fig. 6	N,N'-(propane-1,3-diyl)dinitramide.....	5
Fig. 7	3,7-dinitro-1,3,7-oxadiazocane	5
Fig. 8	5,8-dinitro-1,3,5,8-dioxadiazonane.....	6
Fig. 9	Classic synthesis of DHP	6
Fig. 10	Ring contraction of 2	7
Fig. 11	Ring strain.....	7
Fig. 12	Future targets	8

List of Tables

Table	Performance predictions from Cheetah 7.0	2
-------	--	---

Acknowledgments

The authors wish to thank Terry Piatt and Lori Pridgeon for general equipment assistance and supply management. The authors also wish to thank Eric Bukowski, Brian Roos, and Thuvan Piehler for procedural input and technical review. The authors are indebted to Betsy Rice and Edward Byrd for making available their computer codes and related scripts.

INTENTIONALLY LEFT BLANK.

1. Introduction

Trinitrotoluene (TNT) **3** is a widely used energetic material and was first reported in 1863. By the First World War, all major combatants employed TNT as their most heavily used explosive due to its low melting point, stability, noncorrosive properties, and good performance. Its use predominated in the Second World War in spite of the development of materials such as pentaerythritol tetranitrate, 1,3,5-Trinitroperhydro-1,3,5-triazine, and octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (better known, respectively, as PETN, RDX, and HMX) due to TNT's high productivity. Its use continued through the Vietnam conflict but due to numerous accidents experienced by both the US Navy at sea and the US Air Force in Thailand, alternate binders were sought. The most notable examples were cast-cure, nitramine-based plastic-bonded explosive (PBXs), developed in the 1970s. Cast-cure PBXs, however, are difficult to produce because of incompatibility with existing plants, due to the requirement for high-shear mixers and elevated temperature. Modern alternatives to TNT include eutectic mixtures such as the insensitive munitions explosive IMX-101 and eutectic salts such as DEMN that either have toxicity issues or are difficult to process and incorporate into formulations. Therefore, because of its high productivity, low cost, and favorable processing, the use of TNT is still widespread in military applications around the globe. The search for replacements, however, is ongoing. A series of compounds has been discovered: 3,7-dinitro-1,3,7-oxadiazocane **1** and 5,8-dinitro-1,3,5,8-dioxadiazonane **2** composed of ether-linked nitramines, which are predicted to exhibit explosive properties similar to TNT and have not been described in the literature.




Fig. 1 Structures of Compounds 1–3

Compound **1** is an immediate precursor to melt-castable material 1,3-dinitrohexahydropyrimidine (DHP) **5**, as depicted in Fig. 2, which has TNT's performance and similar physical properties. Furthermore, these compounds undergo a ring-contraction reaction when heated to form either a smaller ether-linked nitramine or a methylene-linked nitramine.




Fig. 2 Ring contraction

Due to their usefulness as synthetic intermediates—and the fact they are not well described in the literature—these ether-linked nitramines warranted further investigation.

2. Computational Analysis and Sensitivity

Compounds **1** and **2** were evaluated using quantum mechanical-modeling codes developed at the US Army Research Laboratory.^{1–5} Using these codes, it becomes possible to accurately predict the heat of formation and density of novel energetic materials. With this information in hand, the compounds can then be evaluated using the Cheetah thermochemical codes to give a theoretical detonation performance. The results of these calculations are listed in the Table below.

Table Performance predictions from Cheetah 7.0

Compound	1	2	TNT ^c
ρ^a	1.60 [1.57]	1.62 [1.61]	1.65
ΔH_f (kJ/mol)	–177.5 ^b	–323.0 ^b	–63.2
P_{cj} (GPa)	21.416	21.328	20.022
D_v (km/s)	7.301	7.491	7.180
ΔH_d (kJ/mL)	7.03	7.19	7.00
OB (%)	–77.61	–58.29	–73.96
Impact (cm)	>152.4	>152.4	>152.4
Friction (N)	>360	>360	>360
ESD (J)	>6.25	>6.25	0.625

^aExperimentally determined values theoretical in brackets.

^bPredicted using the methods of Rice.²

^cHeat of formation and density numbers obtained from Cheetah 7.0 database. Density, ρ ; heat of formation, ΔH_f ; Chapman–Jouguet pressure, P_{cj} ; detonation velocity, D_v ; heat of detonation, ΔH_d ; oxygen balance, OB; electrostatic discharge, ESD.

Using the experimentally determined density with the predicted heat of formation, the results of the Cheetah 7.0 calculations on Compounds **1** and **2** indicate the expected performance of these compounds to be approximately 5% greater than TNT's. Interestingly, both **1** and **2** are quite insensitive; this results in no-go's on small-scale impact tests at heights greater than 152.4 cm. Both compounds were also insensitive to friction stimuli beyond 360 N as well as electrostatic stimuli of greater than 6.25 J (as shown in the Table above).

3. Synthesis Discussion

During this investigation it was discovered that the method of addition of 1,3-dinitramopropane **4** to an acidic formaldehyde solution results either in the direct formation of ether-linked dinitramine **1** or mixtures of **1** and 6-membered dinitramine **5**. (See Fig. 3.)




Fig. 3 Synthesis of Compound **1**

When **4** is added to the reaction mixture as a solution in various solvents, primarily Compound **1** is formed. However, when **4** is added as a pure solid, a mixture of **1** and an impurity that is most likely **5** can be obtained. The best results were produced from very slow addition of **4** dissolved in water, which produced **1** in good yield.

Similarly, when ethylenedinitramine **6** is used in place of **4**, 9-membered heterocycle **2** is obtained (Fig. 4).




Fig. 4 Synthesis of Compound **2**

Both reactions have been performed on up to the 1-g scale. Thermal analysis of both compounds indicates that a decomposition reaction is taking place over a broad range of temperatures starting at 100 °C, which warrants further investigation into the nature of this reaction. The decomposition is accompanied with gas evolution and phase change and is endothermic, according to analysis with differential scanning calorimetry.

4. Results and Experimental Section

The values for the drop-weight experiments were determined using the Langlie one-shot method on an apparatus employing a 2-kg weight from a maximum height of 152.4 cm.⁶ Friction sensitivity measurements were determined on a BAM friction apparatus and ESD was determined using an Allegheny Ballistics Laboratory ESD

apparatus (Table). Nuclear magnetic resonance (NMR) experiments were accomplished using an Anasazi Instruments 90-megahertz (MHz) NMR device. Fourier transform infrared spectroscopy (FTIR) was obtained from a Brüker Alpha-T fitted with a 5-cm path length gas-flow cell in the transmission module, or a diamond-attenuated total-reflectance (DATR) module. Differential scanning calorimetry was conducted on a TA instruments Q10 or Q20 calorimeter. All chemicals were obtained from Sigma-Aldrich, St Louis, Missouri, and were used as received. Values for pH were determined using colorpHast pH-indicator strips (pH 0–14) from EMD Chemicals Inc.

Note: While these compounds were prepared without incident according to the following procedures, these materials are energetic and should be prepared and handled cautiously only by trained personnel.

4.1 Synthetic Procedures

The 2-imidazolidone, available from Sigma-Aldrich, is used to prepare ethylene dinitramine **6** according to the literature method.³

1,3-dinitrotetrahydropyrimidin-2(1H)-one: Was prepared according to a modified literature procedure⁹ in which 1.00 g (10.0 mmole) of N,N' trimethylene urea is added to 3.0 mL of acetic anhydride while stirring in an ice bath. Then, 100% nitric acid (1.5 mL) was slowly added drop-wise over approximately 10 min as to maintain the solution's temperature below 25 °C. Upon addition of 1 mL of the nitric acid, the suspension turned into a solution whereupon the product crashed out as a yellow solid by the time all of the nitric acid was added. Upon completion of nitric-acid addition, the thick suspension was stirred for an additional 20 min, then drowned in ice water. The yellow solid was filtered by vacuum filtration, rinsed with copious amounts of water, and dried under vacuum—or carried immediately to the next step. A typical procedure yields 1.69 g of product (8.85 mmole, 89% yield) after drying in vacuum. (See Fig. 5.)




Fig. 5 1,3-dinitrotetrahydropyrimidin-2(1H)-one

N,N'-(propane-1,3-diyl)dinitramide **4**: Was prepared according to the literature procedure¹⁰ in which 1,3-dinitrotetrahydropyrimidin-2(1H)-one is added to water with stirring and reacted with solid pellets of sodium hydroxide until all solid starting material is dissolved and the pH is strongly alkaline (14 as indicated by pH

paper). Concentrated aqueous hydrochloric acid was added, at which point a large amount of bubbling occurred as carbon dioxide escapes from the reaction. Acid is added until the pH is strongly acidic (0 as indicated by pH paper) and either extraction with ethyl acetate or overnight crystallization in the freezer yielded the title compound as a white solid. (See Fig. 6.)




Fig. 6 N,N'-(propane-1,3-diyl)dinitramide

3,7-dinitro-1,3,7-oxadiazocane **1**: Was prepared by the dropwise addition of 1.0 g (6.0 mmole) of N,N'-(propane-1,3-diyl)dinitramide in 10 mL of water to 22 mL of a 73% sulfuric-acid–water solution containing 1.4 g (46 mmole) of paraformaldehyde. The addition rate was controlled such as to keep the solution's temperature below 7 °C (addition occurred over 15 min). After the addition was completed the solution was stirred for an additional 5 min whereupon it turned milky with a white precipitate, which was drowned in ice water and filtered to yield 796 mg (3.86 mmole, 64%) of **1** as a white solid after drying under vacuum. ¹H NMR (90.420 MHz, Acetone-D₆) δ 5.40 (s, 4H), δ 4.06 (dd, 4H, j = 5.15 Hz, 5.15 Hz), δ 1.98 (m, 2H); ¹³C NMR (22.605 MHz, Acetone-D₆) δ 81.74 (2C), 52.00 (2C), 24.27 FTIR (DATR), $\tilde{\nu}$ = 3064, 1518, 1386, 1286, 1254, 1234, 995, 959, 874, 762, 583. (See Fig. 7.)




Fig. 7 3,7-dinitro-1,3,7-oxadiazocane

5,8-dinitro-1,3,5,8-dioxadiazonane **2**: Was prepared by the dropwise addition of 1.00 g (6.67 mmole) of N,N'-(ethane-1,2-diyl)dinitramide dissolved in 8.5 mL 1,4-dioxane to 20 mL of a 73% sulfuric-acid–water solution containing 1.40 g (46 mmole) of paraformaldehyde. The nitramine solution was prepared by heating the dioxane until the nitramine dissolved. This solution was allowed to cool but not to the point of precipitation. The warm nitramine solution was added to the acidic formaldehyde solution before precipitation of the nitramine occurred. The sulfuric acid solution was chilled down to below 5 °C prior to addition of the nitramine solution. The addition was completed in about 30 s using a pipet to add the solution nitramine solution to the chilled acidic formaldehyde solution. After the addition was completed the solution was stirred for an additional 5 min whereupon it turned milky with a white precipitate, which was drowned in ice water and filtered to yield

1.05 g (4.77 mmole, 72%) of **1** as a white solid after drying under vacuum. ^1H NMR (90.420 MHz, Acetonitrile- D_3) δ 5.40 (s, 4H), δ 5.03 (s, 2H), δ 4.16 (s, 4H); ^{13}C NMR (22.605 MHz, Acetonitrile- D_3) δ 99.24, 79.92 (2C), 47.377 (2C); FTIR (DATR), $\tilde{\nu}$ = 3040, 2966, 1511, 1458, 1264, 1264, 1101, 1003, 989. (See Fig. 8.)




Fig. 8 5,8-dinitro-1,3,5,8-dioxadiazonane

5. Discussion of DHP and Formaldehyde Extrusion from Ether-Linked Nitramines

DHP has been synthesized by a classic ring closure of the primary nitramine precursor (Fig. 9).





Fig. 9 Classic synthesis of DHP

The classic synthesis has a low overall yield and requires a very large amount of sulfuric acid, generating excess acidic waste. The synthesis of **1** as a precursor to DHP is much higher yielding and allows a substantial reduction in the amount of sulfuric acid used. Compound **1** undergoes thermal ring contraction with extrusion of a molecule of formaldehyde to efficiently form DHP.

Compound **2** also undergoes thermal-ring contraction with extrusion of formaldehyde to form Compound **7** but does not contract down to the 5-membered nitramine **8** (Scheme 5), as depicted in Fig. 10.



It is speculated that ring strain may account for the failure to undergo thermally induced ring contraction from the **7** to **8**, while the 8-membered ring readily undergoes this transformation (Fig. 11).




Fig. 11 **Ring strain**

Further development will be focused on increasing the scope of the ring contraction by screening solvents and various acid catalysts to promote the described transformation. Preliminary results suggest this will be possible because traces of **8** have been detected by gas chromatography–mass spectrometry in small-scale screening reactions which utilizing nitromethane as a solvent and catalyst, which cause **7** to undergo reactions. While mostly decomposition has been observed, it is thought that with more screening, the reaction can be driven to a more productive pathway.




Fig. 12 Future targets

Compound **9** has been published in the Russian literature as having a melting point of 100 °C, but the synthesis has been unreproducible. Compound **10** is a potential precursor to **11**, which is a more stable analog of **12**, a very powerful but moisture-sensitive explosive. However, **11**, lacking carbonyl groups, is expected not to exhibit moisture sensitivity but retain much of the performance of **12**. It is thought that through careful screening of conditions, **10** can be synthesized and forced to undergo ring contraction to form **11**.

7. References

1. Byrd EFC, Rice BM. Improved prediction of heats of formation of energetic materials using quantum mechanical calculations. *J Phys Chem A*. 2006;110:1005–1013.
2. Rice BM, Pai SV, Hare J. Predicting heats of formation of energetic materials using quantum mechanical calculations. *Combust Flame*. 1999;118:445–458.
3. Rice BM, Hare JJ, Byrd EFC. Accurate predictions of crystal densities using quantum mechanical molecular volumes. *J Phys Chem A*. 2007;111:10874.
4. Byrd EFC, Rice BM. A comparison of methods to predict solid phase heats of formation of molecular energetic salts. *J Phys Chem A*. 2009;113:345.
5. Rice BM, Byrd EFC. Evaluation of electrostatic descriptors for predicting crystalline density. *J Comput Chem*. 2013;34:2146–2151.
6. Langlie HJ. A reliability test for “one-shot” items. Technical Report U-1792 Aeronutronic Division of Ford Motor Company. Newport Beach (CA): 1965.
7. Bastea S, Fried LE, Glaesemann KR, Howard WM, Kuo IW, Souers PC, Vitello PA. Cheetah 7.0 user’s manual. Livermore (CA): Lawrence Livermore National Laboratory; 2012. Report No.: LLNL-SM-599073.
8. Bachmann WE, Horton WJ, Jenner EL, MacNaughton NW, Maxwell CE. *J Am Chem Soc*. 1950;72:3132.
9. McKay AF, Wright GF. The nitration products of 2-nitramino- Δ^2 -1,3-diazacycloalkenes. *J Am Chem Soc*. 1948;70:3990.
10. McKay AF, Manchester DF. The nitration products of some substituted 2-nitramino-1,3-diazacycloalkenes. *J Am Chem Soc*. 1949;71:1970.

List of Symbols, Abbreviations, and Acronyms

ATR	attenuated total reflectance
cm	centimeter
CO ₂	carbon dioxide
DATR	diamond-attenuated total-reflectance
DHP	1,3-dinitrohexahydropyrimidine
D _v	detonation velocity
ESD	electrostatic discharge
FTIR	Fourier transform infrared spectroscopy
g	gram
GPa	gigapascal
ΔH _d	heat of detonation
ΔH _f	heat of formation
HCl	hydrochloric acid
HNO ₃	nitric acid
H ₂ SO ₄	sulfuric acid
H ₅₀	50% initiation height
IMX	insensitive munitions explosive
J	joule
kJ	kilojoule
MHz	megahertz
min	minute
mL	milliliter
mmole	millimole
mol	mole
N	newton
N ₂	nitrogen
NaOH	sodium hydroxide
NMR	nuclear magnetic resonance

OB	oxygen balance
PBX	plastic-bonded explosive
P_{cj}	Chapman–Jouguet pressure
ρ	density
s	second
TNT	trinitrotoluene
ZnSe	Zinc Selenide

1 (PDF)	DEFENSE TECHNICAL INFORMATION CTR DTIC OCA	RDRL WML C S AUBERT J BANNING K MCNESBY
2 (PDF)	DIRECTOR US ARMY RESEARCH LAB RDRL CIO L IMAL HRA MAIL & RECORDS MGMT	B ROOS RDRL WML H J NEWILL
1 (PDF)	GOVT PRINTG OFC A MALHOTRA	
2 (PDF)	WEAPONS DEV & INTEGRATION DIRCTRT AMRDEC RDMR WDN J NEIDERT G DRAKE	
1 (PDF)	US ARMY ARDEC AMSRD AAR AEE W R DAMAVARAPU	
1 (PDF)	US ARMY ARDEC RDAR MEE W K SPANGLER	
1 (PDF)	LOS ALAMOS NATL LAB D CHAVEZ	
1 (PDF)	LAWRENCE LIVERMORE NATL LAB P PAGORIA	
2 (PDF)	NAVAL SURFACE WARFARE CTR B SLEADD P CARUANA	
13 (PDF)	DIR USARL RDRL CI P PLOSTINS RDRL SL P BAKER RDRL WM B FORCH J ZABINSKI RDRL WML M ZOLTOSKI RDRL WML B E BYRD B RICE N TRIVEDI	